

A & Z - Children's Aspirin (NSAID) Orange Flavored Chewable Tablets, 81 mg

Drug Facts

Active Ingredient (in each tablet)

Aspirin 81 mg (NSAID*).....Analgesic/Antipyretic
*non-steroidal anti-inflammatory drug

Purpose

Uses ■ temporary relief of minor aches, pains and headaches ■ to reduce fever associated with colds, sore throats, teething

Warnings

Reye's Syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's Syndrome, a rare but serious illness.

Allergy Alert : Aspirin may cause a severe allergic reaction which may include hives, facial swelling, asthma (wheezing), shock.

Stomach Bleeding Warning: This product contains a non-steroidal anti-inflammatory drug (NSAID), which may cause stomach bleeding. The chance is higher if the user ■ has had stomach ulcers or bleeding problems ■ takes a blood thinning (anticoagulant) or steroid drug ■ takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ takes more or for a longer time than directed ■ is age 60 or older ■ has 3 or more alcoholic drinks every day while using this product

Sore Throat Warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly

Do not use ■ if you are allergic to aspirin or other pain relievers/fever reducers ■ for at least 7 days after tonsillectomy or oral surgery

Ask a doctor before use if ■ stomach bleeding warning applies to user ■ user has history of stomach problems, such as heartburn ■ user has high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ user is taking a diuretic ■ user has not been drinking fluids ■ user has lost a lot of fluids due to vomiting or diarrhea

Ask a doctor or pharmacist before use if you are taking a prescription drug for ■ anticoagulation (thinning the blood) ■ diabetes ■ gout ■ arthritis

Stop use and ask a doctor if ■ user experiences any of the following signs of stomach bleeding ■ feels faint, vomits blood, has bloody or black stools, has stomach pain that does not get better ■ an allergic reaction occurs. Seek medical help right away ■ new symptoms occur ■ redness or swelling is present ■ ringing in the ears or loss of hearing occurs ■ pain gets worse or lasts for more than 5 days ■ fever gets worse or lasts for more than 3 days ■ **You may report side effects to 1-800-952-0050**

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery

Keep out of reach of children. In case of accidental overdose, contact a doctor or Poison Control Center immediately.

Directions to be administered only under adult supervision ■ repeat every 4 hours, while symptoms persist, up to a maximum of five doses in 24 hours or as directed by a doctor ■ drink water with each dose ■ Children under 3 years: consult a doctor

Age (yrs)	3 to under 4	4 to under 6	6 to under 9	9 to under 11	11 to under 12	Adults and children 12 years and over
Weight (lbs)	32 to 35	36 to 45	46 to 65	66 to 76	77 to 83	
Dosage	2 tablets	3 tablets	4 tablets	4 – 5 tablets	4 – 6 tablets	5 – 8 tablets

Other Information ■ store at room temperature

Inactive Ingredients flavor, FD&C yellow # 6, saccharin sodium, silicon dioxide, starch, stearic acid, sucrose